

Contribution to the study of the ground-beetle fauna (Coleoptera: Carabidae) of the Osogovo Mountain, Bulgaria. III.

Borislav GUÉORGUIEV

Introduction

In the first part of the present study the author adds 77 more species and subspecies which are new for the carabid fauna of the Osogovo Mt, as well as data about their vertical range, season activity and life conditions of the adults. All specimens have been collected from the Bulgarian part of the mountain. The second half of the paper concerns the erroneous identifications made in all the previous papers on this subject, and the determination of the subspecies status of the ground-beetles from the region.

Material and methods

The bulk of the specimens were collected by the author during the period of April - November 1994, April - October 1995, April - October 1996, October 1997 and June - July 1998. Single specimens collected by other colleagues before 1994 have been enlisted too. The collection methods are the same as indicated by GUÉORGUIEV (1996). The material is preserved in the National Museum of Natural History, Sofia (NMNH). The habitats visited by the author are:

H 1. Hisarluka Park, 640-670 m, below Bogoslov Village. Secondary coniferous plantations (*Pinus nigra* predominating, as well as *P. silvestris* and *Corylus avellana*).

H 2. East slope above Bogoslov Village, 900 m. Meadows.

H 3. North slope above the road Bogoslov Village - Trite Bouki Chalet, 940-960 m. Beech forest.

H 4. Popovi Livadi, 1230-1260 m. Beech forest.

H 5. Popovi Livadi, 1230-1260 m. Meadows used for pastures.

- H 6.** Iglika Chalet, 1320-1340 m. Spruce forest.
- H 7.** Iglika Chalet, 1330-1350 m. Beech forest.
- H 8.** Iglika Chalet, 1340-1350 m. Meadows. Moist habitat around brook.
- H 9.** Trite Bouki Chalet and Bor Chalet, 1540-1570 m. Beech forest.
- H 10.** Trite Bouki Chalet, 1540-1570 m. Coniferous forest (*Pinus sylvestris*).
- H 11.** Trite Bouki Chalet, 1550-1650 m. Secondary grass vegetation in the place of former beech and coniferous forests.
- H 12.** The section of Trite Bouki Chalet - Begbunar Spring, 1600-1830 m. Grass vegetation. Moist habitats.
- H 13.** Begbunar Spring, 1820-1840 m. Hygrophilous grass vegetation.
- H 14.** The section of Begbunar Spring - the south foot of Tsurni Kamak Peak, 1830-2000 m. The orophytic belt above Gurlyano Village is also includet there, 1800-2000 m. Grass vegetation.
- H 15.** The section of the south foot of Tsurni Kamak Peak - Ruen Peak, 2000-2251 m. The orophytic belt above Gurlyano Village is also includet there, 2000-2251 m. Grass vegetation.
- H 16.** Riverside places before and after Stradalovo Village, 700-720 m. Stony and muddy habitats.
- H 17.** Xerophytic terrain between Pelatikovo Village and Stradalovo Village, 800 m. Secondary coniferous forest.
- H 18.** The Eleshnitsa Valley between Stradalovo Village and Rakovo Village, 800-850 m. Hygrophilous riverside vegetation on calcareous terrain.
- H 19.** Brook between Rakovo Village and Vetren Village, 850 m. Xerophytic vegetation on silicate terrain.
- H 20.** Chekanetski bridge over the Eleshnitsa River, 950 m. Xerophytic grass. Deciduous wood riverside vegetaton.
- H 21.** The Eleshnitsa Valley between Chekanetski bridge and Sazhdenik Village, 1000-1050 m. Grass and deciduous wood riverside vegetaton.
- H 22.** Hygrophilous to hydrophilous stony place near the Eleshnitsa Valley, 1300 m. Around a beech forest.
- H 23.** North slope, south of the Eleshnitsa Valley, 1350 m. Beech forest.
- H 24.** The Eleshnitsa Valley, 1350 m. Riverside meadow.
- H 25.** Mechata Doupka Cave near Stradalovo Village, 850 m. Xerophytic oak forest on calcareous terrain.
- H 26.** Along the Novoselska River between Slokoshtitsa Village and Novo Selo Village, 720 m. Slimy rush habitat.
- H 27.** The place of Dve Reki along the Novoselska River, 1000 m. Various habitats: riverside vegetation (*Populus alba* and *Alnus* spp.); potato fields; coniferous forest.
- H 28.** Big torrent near the road Trite Bouki Chalet - Novo Selo Village, 1300-1350 m. Mixed coniferous and mesophilous deciduous forest.
- H 29.** Hygrophilous to hydrophilous riverside place at the Mlachka Reka River near Chervena Yabulka Village, 1440-1460 m. Mixed forest.

List of the species and subspecies

***Omophron (Omophron) limbatus* (F.).** H 16 (2-3.VII.1998, 2 specimens); H 20 (5.VI.1998, 4 specimens); H 26 (2.VII.1998, 2 specimens).

***Leistus (Pogonophorus) magnicollis magnicollis* Motsch.** H 9 (2.IX.1994, 1 ♀, in beach foliage; 7.VI.1995, 1 ♂, in beech foliage). Balkan endemic species and subspecies.

***Leistus (Leistus) ferrugineus* (L.).** H 5 (traps: V.1996, 1 specimen; IX-X.1996, 1 specimen); H 8 (3.VI.1998, 1 ♀); H 9 (traps: IX-X.1995, 1 ♂).

***Nebria (Nebria) brevicollis* (F.).** H 19 (traps: V.1996, 1 ♂).

***Notiophilus (Notiophilus) germinyi* Fauv.** H 14 (Choveka Peak, 2000 m, 31.VII.1980, 2 specimens, leg. J. Ganev).

***Calosoma (Calosoma) sycophanta* (L.).** H 21 (2.VII.1998, remains of elytrae).

***Elaphrus (Neoelaphrus) uliginosus* F.** H 26 (2.VII.1998, 1 ♀).

***Elaphrus (Elaphroterus) aureus aureus* P. Mull.** H 16 (2-3.VII.1998, 1 ♂, 1 ♀); H 21 (5.VI.1998, 2 ♀♀; 2.VII.1998, 1 ♂).

***Loricera (Loricera) pilicornis pilicornis* (F.).** H 27 (5.VI.1998, 2 ♂♂).

***Dyschiriodes (Dyschiriodes) politus politus* Dej.** H 26 (2.VII.1998, 1 specimen).

***Broscus cephalotes* (L.).** H 16 (2-3.VII.1998, 2 ♂♂).

***Perileptus (Perileptus) areolatus areolatus* (Creutzer).** H 16 (2-3.VII.1998, 8 specimens).

***Trechus (Trechus) quadristriatus* (Schrk.).** H 3 (7.VI.1995, 1 specimen); H 9 (2-3.IX.1996, 2 specimens, in beech foliage); H 16 (2-3.VII.1998, 1 specimen).

***Trechus (Trechus) austriacus* Dej.** H 25 (traps: IX.1994-II.1995, 1 specimen).

***Trechus (Trechus) subnotatus subnotatus* Dej.** H 9 (2.IX.1994, 1 specimen, in beech foliage near a brook; 2.X.1995, 5 specimens, near a brook; 7.VI.1996, near a brook, 2 specimens; 2-3.IX.1996, 1 specimen, in beech foliage); H 11 (3.VI.1998, 1 specimen); H 21 (5.VI.1998, 1 specimen); H 22 (traps: VI-VII.1996, 1 specimen).

***Tachys (Paratachys) bistriatus* (Duft.).** H 16 (2-3.VII.1998, 2 specimens).

***Elaphropus (Tachyura) quadrisignatus* (Duft.).** H 14 (Tsurni Kamak Peak, 2000 m, 5.VII.1995, 3 specimens); H 21 (9-10.VI.1995, 2 specimens).

***Elaphropus (Tachyura) diabrychys* (Kol.).** H 16 (2-3.VII.1998, 8 specimens); H 21 (2.VII.1998, 2 specimens); H 27 (9.VI.1995, 1 specimen; 5.VI.1998, 1 specimen).

***Tachyta nana nana* (Gyllh.).** H 11 (3.VI.1998, 2 specimens under the bark of a beech log).

***Bembidion (Metallina) lampros* (Hbst.).** in large numbers in: H 3 (27.IV.1994); H 4 (10.VII.1994); H 8 (25.IV.1995); H 9 (2.X.1995); H 11 (3.VI.1998);

H 12 (4.VI.1994; 8.VI.1995; 4.VI.1998); H 13 (4.VI.1994; 8.VI.1995; 4.VI.1998); H 15 (Shapka Peak, 2150 m, 8.VI.1995, snow spots); H 18 (11.V.1996); H 21 (9-10.VI.1995; 5.VI.1998; 2.VII.1998); H 28 (5.IV.1994); H 29 (2.IX.1994; 5.VIII.1996).

***Bembidion (Metallina) properans* Steph.** H 5 (traps: V.1994, 2 ♂♂, 2 ♀♀; V.1996, 2 ♀♀); H 8 (3.VI.1998, 2 specimens); H 12 (8.VI.1995, 4 specimens); H 27 (2.VII.1998, 1 specimen).

***Bembidion (Synechostictus) millerianum* Heyd.** H 14 (Tsurni Kamak Peak, 2000 m, 5.VII.1995, 1 specimen); H 20 (5.VI.1998, 1 specimen); H 21 (27-28.IV.1994, 1 specimen; 9-10.VI.1995, in large numbers; 5.VI.1998, in large numbers); H 27 (5.VI.1998, 1 specimen).

***Bembidion (Nepha) caucasicum* Motsch.** H 12 (4.VI.1994, 2 specimens; 4.VI.1998, 1 specimen); H 13 (4.VI.1994, 7 specimens; 4.IX.1995, 2 specimens); H 14 (Tsurni Kamak Peak, 1950-2000 m, 4.VI.1994, 5 specimens; 16.X.1997, 1 specimen; traps: VII.1994, 3 specimens); H 15 (Shapka Peak, 2000-2100 m, 8.VI.1995, 1 specimen, snow spots).

***Bembidion (Nepha) tetragrammum illigeri* Net.** H 16 (2-3.VII.1998, 1 ♂).

***Bembidion (Bembidionetolitzkya) varicolor varicolor* (F.).** H 16 (2-3.VII.1998, 1 ♂, 2 ♀♀).

***Bembidion (Bembidionetolitzkya) rhodopense* Apf.** H 12 (1.IX.1994, 1 specimen); H 27 (9.VI.1995, 1 specimen, near a brook). Balkan endemic species.

***Bembidion (Bembidionetolitzkya) tibiale* (Duft.).** H 16 (2-3.VII.1998, 1 ♂, 1 ♀); H 21 (5.VI.1998, 1 ♀); H 27 (2.VII.1998, 1 ♀).

***Bembidion (Bembidionetolitzkya) geniculatum geniculatum* Heer.** H 14 (16.X.1996, 1 ♀).

***Bembidion (Peryphanes) deletum deletum* Serv.** H 8 (6.VIII.1996, 1 ♀; 3.VI.1998, 1 ♀); H 12 (4.VI.1994, 2 specimens; 1.IX.1994, 2 specimens; 8.VI.1995, 4 specimens, 22.X.1995, 1 specimen; 4.VI.1998, 2 specimens); H 13 (4.VI.1994, 3 specimens; 8.VI.1995, 1 ♂, 1 ♀; 4.IX.1995, 2 specimens); H 14 (Tsurni Kamak Peak, 2000 m, 4.VI.1994, 2 specimens); H 15 (Shapka Peak, 2150 m, 8.VI.1995, 1 specimen, snow spots); H 29 (2.IX.1994, 1 specimen).

***Bembidion (Peryphanes) dalmatinum dalmatinum* Dej.** H 9 (2.IX.1994, 1 specimen); H 26 (2.VII.1998, 1 ♂); H 27 (5.VI.1998, 1 specimen).

***Bembidion (Peryphanes) pindicum* Apfelbeck, 1901.** H 14 (the south foot of Tsurni Kamak Peak, 2000 m, 16.X.1997, 1 ♀, leg. B. Guéorguiev, under stone in the stratum). **New for the Bulgarian fauna.** A Balkan endemic species. According to APFELBECK (1902; 1904) this taxon is distributed in the alpine regions of the following high Balkan mountains: Taygetos Mt (2407 m) in Pelopponesos; Timfristos (= Veluchi) Mt (2315 m), the massifs of Oxia (1926 m) and Korax (2350 m) - both parts of Vardousia Mt, Parnassos Mt (2457 m) - all situated in Central Greece; Peristeri (2295 m) - a massif of Pindos Mts in north-western Greece; further north on the Vran Mt (2074 m) in south Bosnia. Probably a more peculiar form, something intermediate between *B. pindicum* and *B. asi-*

aeminoris Net., inhabits the Korabi Mt (2764 m) at the Albanian-Macedonian border (NETOLITZKY, 1943). Its occurrence on Osogovo Mt (2251 m), on the east of the Vardar zoogeographical barrier, is quite interesting.

***Bembidion (Peryphanes) stephensi stephensi* Crotch.** H 29 (5.VIII.1996, 1 ♀).

***Bembidion (Peryphanes) grandipenne grandipenne* Schaum.** H 26 (2.VII.1998, 1 ♀).

***Bembidion (Peryphus) femoratum femoratum* Sturm.** H 16 (2-3.VII.1998, 1 ♂); H 26 (2.VII.1998, 1 ♀).

Bembidion (Peryphus) subcostatum javurkovaе Fass. in large numbers in: H 11 (3.VI.1998); H 16 (2-3.VII.1998); H 20 (27-28.IV.1994; 5.VI.1998; traps: V.1994); H 21 (9-10.VI.1995; 5.VI.1998); H 27 (9.VI.1995; 5.VI.1998); H 28 (5.IV.1994; 9.VI.1995); H 29 (2.IX.1994).

***Bembidion (Ocydromus) decorum decorum* (Zenker).** H 16 (2-3.VII.1998, 1 ♂, 2 ♀♀).

***Pterostichus (Phonias) strenuus* (Panz.).** H 20 (traps: V.1996, 1 ♀).

***Pterostichus (Argutor) vernalis* (Panz.).** H 8 (traps: VI-VII.1996, 1 ♀).

***Pterostichus (Haptoderus) brevis brevis* (Duft.).** H 6 (traps: IX-X.1995, 1 ♂; IX-X.1996, 1 ♀); H 7 (traps: V.1996, 1 ♂, 1 ♀; VI-VII.1996, 1 ♂); H 9 (traps: VI-VII.1996, 2 ♀♀; VI-VII.1996, 1 ♂); H 10 (traps: V.1996, 1 ♀; VIII.1996, 1 ♀); H 22 (traps: VI-VII.1996, 1 ♂, 1 ♀; VIII.1996, 1 ♀). So far known in Bulgaria only from Belasitsa Mt (GUÉORGUIEV, 1992).

***Pterostichus (Haptoderus) vecors* Tschitsch.** H 18 (traps: V.1996 2 ♂♂); H 22 (5.V.1996, 1 ♂, 1 ♀; traps: VI-VII.1996, 1 ♂). A Bulgarian endemic species.

***Pterostichus (Morphnosoma) melanarius bulgaricus* Lutsh.** H 27 (5.VI.1998, 1 ♂).

***Pterostichus (Pterostichus) brucki* Schaum.** in large numbers in: H 2 (5.VIII.1996); H 6 (traps: VI-VII.1996; VIII.1996); H 7 (traps: V.1996; VI-VII.1996); H 9 (traps, V.1996; VI-VII.1996; VIII.1996); H 18 (traps: V.1996; VI-VII.1996; VIII.1996); H 22 (traps: V.1996; VI-VII.1996; VIII.1996); H 23 (traps: VI-VII.1996; VIII.1996); H 24 (5.V.1996; traps: VI-VII.1996; VIII.1996); H 27 (6.VIII.1996); H 29 (5.VIII.1996).

***Olistopus sturmi* (Duft.).** H 5 (traps: IV.1994, 1 specimen).

***Agonum (Agonum) duftschmidi* Schmidt.** M 14.(8.VI.1995, 2 ♀♀).

***Agonum (Agonum) muelleri* (Hbst.).** H 26 (2.VII.1998, 1 ♂).

***Agonum (Agonum) viridicupreum viridicupreum* (Goeze).** M 8 (3.VI.1998, 1 ♀).

***Synuchus (Synuchus) vivalis vivalis* (Ill.).** H 3 (traps: VI-VII.1996, 1 ♂); H 20 (traps: VI-VII.1996, 1 ♂, 1 ♀); H 21 (9-10.VI.1995, 1 ♂).

***Amara (Amara) littorea* Thoms.** H 7 (traps: V.1996, 1 ♀); H 9 (4.V.1996, 1 ♂); H 24 (traps: V.1996, 1 ♂).

***Amara (Amara) proxima* Putz.** H 11 (8.VI.1995, 1 specimen).

- Amara (Amara) anthobia*** Villa. H 18 (traps: V.1996, 1 ♂).
Amara (Amara) lucida (Duft.). H 18 (11.V.1996, 1 ♂).
Amara (Celia) ingenua (Duft.). H 12 (22.X.1995, 4 specimens).
Amara (Bradytus) fulva (O. F. Muller). H 27 (2.VII.1998, 1 ♀).
Curtonotus (Curtonotus) aulicus (Panz.). H 18 (traps: VI- VII.1996, 1 ♀); H 24 (traps: VIII.1996, 2 ♂♂, 2 ♀♀).
Anisodactylus (Anisodactylus) binotatus (F.). H 26 (2.VII.1998, 1 ♂).
Stenolophus (Stenolophus) teutonus (Schrk.). H 26 (2.VII.1998, 4 ♀♀); H 27 (5.VI.1998, 1 ♂, 1 ♀; 2.VII.1998, 2 ♀♀).
Acupalpus (Acupalpus) flavicollis (Sturm). H 26 (2.VII.1998, 2 specimens); H 27 (5.VI.1998, 7 specimens; 2.VII.1998, 1 specimen).
Harpalus (Pseudophonus) griseus (Panz.). H 27 (5.VI.1998, 1 ♂).
Harpalus (Harpalus) honestus (Duft.). H 21 (9-10.VI.1995, 1 ♂).
Harpalus (Harpalus) sulphuripes sulphuripes Germ. H 2 (10.V.1996, 1 ♂, 2 ♀♀); H 8 (traps: V.1996, 1 ♀).
Harpalus (Harpalus) attenuatus Steph. H 2 (traps: V.1996, 1 ♂); H 5 (traps: IX-X.1995, 1 ♀); H 19 (traps: VI-VII.1996, 1 ♀).
Harpalus (Harpalus) triseriatus triseriatus Fleisch. H 17 (10.VI.1995, 1 ♂).
Harpalus (Harpalus) tardus (Panz.). H 1 (traps: V.1996, 1 ♂, 1 ♀); H 18 (traps: V.1996, 9 ♂♂, 2 ♀♀; VI-VII.1996, 5 ♂♂, 8 ♀♀).
Harpalus (Harpalus) latus (L.). H 8 (traps: V.1996, 1 ♂, 1 ♀; 3.VI.1998, 1 ♂); H 27 (5.VI.1998, 1 ♂).
Harpalus (Harpalus) smaragdinus (Duft.). H 5 (traps: VI-VII.1996, 1 ♂; VIII.1996, 1 ♀); H 14 (below Shapka Peak, 2000 m, 8.VI.1995, 1 ♂, 1 ♀); H 19 (traps: VI-VII.1996, 1 ♀); H 26 (2.VII.1998, 1 ♀).
Ophonus (Metophonus) puncticeps Steph. H 18 (traps: VIII.1996, 1 ♀); H 22 (traps: VI-VII.1996, 1 ♀).
Ophonus (Metophonus) schaubergerianus Puel. H 18 (traps: V.1996, 1 ♀); H 20 (traps: V.1996, 2 ♂♂, 1 ♀); H 21 (2.VII.1998, 1 ♀); H 24 (traps: VIII.1996, 1 ♂).
Ophonus (Hesperophonus) azureus (F.). H 5 (traps: V.1996, 1 ♂); H 8 (25.IV.1995, 1 ♀); H 16 (2-3.VII.1998, 1 ♂).
Ophonus (Ophonus) sabulicola ponticus Schaub. H 5 (traps: IX-X.1995, 1 ♂).
Panagaeus (Panagaeus) bipustulatus (F.). H 20 (traps: V.1995, 1 ♂); H 21 (9-10.VI.1995, 1 ♂).
Chlaenius (Chlaeniellus) nitidulus (Schrk.). H 16 (2-3.VII.1998, 1 ♂, 1 ♀); H 21 (9-10.VI.1995, 1 ♂; 5.VI.1998, 1 ♀); H 26 (2.VII.1998, 2 ♀♀); H 27 (5.VI.1998, 1 ♂, 1 ♀; 2.VII.1998, 1 ♀).
Chlaenius (Chlaeniellus) vestitus (Payk.). H 8 (6.VIII.1996, 1 ♀).
Syntomus pallipes (Dej.). H 2 (10.V.96, 1 specimen); H 8 (23.X.1995, 1 spec-

imen); H 20 (5.VI.1994, 2 specimens; 5.V.1996, 1 specimen); H 21 (28.IV.1994, 1 specimen; 9-10.VI.1995, 1 specimen); H 27 (9.VI.1995, 1 specimen).

***Syntomus truncatellus truncatellus* (L.).** H 8 (25.IV.1995, 1 specimen); H 12 (8.VI.1995, 2 specimens); H 14 (16.X.1997, 1 specimen); H 26 (2.VII.1998, 1 specimen).

***Microlestes fissuralis* Rtt.** H 8 (25.IV.1995, 1 specimen).

***Lionychus quadrillum* (Duft.).** H 16 (2-3.VII.1998, 3 ♂♂, 4 ♀♀); H 27 (5.VI.1998, 1 ♂).

***Cymindis (Cymindis) lineata* (Quens.).** H 5 (traps: VI.1995, 1 ♂).

Another ground-beetle - *Harpalus (Harpalus) serripes serripes* (Quens.), found in the Macedonian part of the Osogovo Mt (GUÉORGUIEV, 1996) has been also established in the Bulgarian one - H 5 (traps: VI.1995, 1 ♂); H 18 (11.V.1996, 1 ♂, 1 ♀); H 21 (27.IV.1995, 1 ♂; 9-10.VI.1995, 1 ♂; 5.VI.1998, 1 ♂); H 26 (2.VII.1998, 1 ♀).

Critical notes on published taxa of ground-beetles from Osogovo Mt

Some taxa published by BURESCH & KANTARDJIEVA (1928), DRENSKY (1928), KANTARDJIEVA (1928), PAWLOWSKI (1973), HIEKE & WRASE (1988), and GUÉORGUIEV (1996) have either been determinated wrongly or the subspecific status of the local populations has not been specified correctly. After a revision, done by Dr Hieke, Mr. Wrase and the present author, their correct identification is indicated below.

***Cicindela (Cicindela) campestris campestris* L.**, det. Guéorguiev (KANTARDJIEVA, 1928, sub *C. c.* var. *palustris* Motsh.).

***Carabus (Carabus) ullrichi fastuosus* Pall.**, det. Guéorguiev (BURESCH & KANTARDJIEVA, 1928, sub *C. (Eucarabus) u. rhilensis* Kr.).

***Carabus (Tomocarabus) convexus dilatatus* Dej.**, det. Guéorguiev (GUÉORGUIEV, 1996, sub *C. (Morphocarabus) scabriusculus bulgarus* Lap.). This taxon has been cited from Osogovo Mt (BURESCH & KANTARDJIEVA, 1928).

***Carabus (Archicarabus) montivagus montivagus* Pall.**, det. Guéorguiev (BURESCH & KANTARDJIEVA, 1928, sub *C. (Deuterocarabus) m.* Pall.; GUÉORGUIEV, 1996, sub *C. (A.) m. bulgaricus* Csiki).

***Carabus (Pachystus) cavernosus cavernosus* Friv.**, det. Guéorguiev (BURESCH & KANTARDJIEVA, 1928, sub *C. (P.) c.* Friv.; DRENSKY, 1928, sub *C. c.* Friv.; HIEKE & WRASE, 1988, sub *C. (P.) c.* Friv.).

***Carabus (Chaetocarabus) intricatus intricatus* L.**, det. Guéorguiev (BURESCH & KANTARDJIEVA, 1928, sub *C. (Ch.) i. starensis* Born.).

***Clivina collaris collaris* (Hbst.)**, det. Wrase (GUÉORGUIEV, 1996, sub *C. fessor* (L.)). First record for Osogovo Mt.

***Trechus priapus medius* Meixner**, det. Guéorguiev (PAWLOWSKI, 1973, sub

Tr. p. K. Dan.). Sciaky (in litt.) determined the local specimens first as *T. priapus*. Then it was established from the almost unknown paper of MEIXNER (1939) that a separate geographical race of that species - *T. p. medius*, was described which is a subspecies inhabiting the area between Morava River and Iskar River.

Platynus (Platynidius) scrobiculatus serbicus (Csiki), det. Guéorguiev (GUÉORGUIEV, 1996, sub *Agonum (Platynus) s. F.*).

Calathus (Calathus) fuscipes fuscipes (Goeze), det. Guéorguiev (GUÉORGUIEV, 1996, sub *C. (C.) f. (Goeze)*). After the publication of the last paper of the author, it was established that the material published as *C. (C.) f.* consists of the two close taxa - *C. fuscipes fuscipes* and *C. distinguendus Chd.* Using the same habitats (H) given in GUÉORGUIEV (1996), the data pertaining to *C. (C.) distinguendus* are presented below. The remaining information concerns *C. f. fuscipes*.

Calathus (Calathus) distinguendus Chd., det. Guéorguiev. Tash-Tepe Peak, 1996 m, 21.VI.1926, 2 ♂♂, 2 ♀♀; H 1 (traps: V., 1 ♀; VII., 2 ♀♀; VIII., 1 ♂, 2 ♀♀; IX., 1 ♀); H 2 (27.IV., 1 ♂); H 6 (VII.-IX., 2 ♂♂, 2 ♀♀); H 8 (traps: V., 1 ♂; VII., 1 ♀; VIII., 1 ♀; IX., 2 ♀♀); H 9 (25.III., 1 ♀; traps: IV., 1 ♂, 3 ♀♀; VI., 1 ♂; VII., 2 ♂♂, 2 ♀♀; VIII., 2 ♂♂, 3 ♀♀; IX., 15 ♂♂, 31 ♀♀); H 19 (5.VI.1998, 1 ♂). A new taxon for Osogovo Mt.

Amara (Amara) nitida nitida Sturm, det. Guéorguiev (HIEKE & WRASE, 1988, sub *A. (A.) n. Sturm*).

Amara (Amara) lunicollis Schdt., det. Hieke (GUÉORGUIEV, 1996, sub *A. (A.) morio nivium Tschitsch.*). A first record for Osogovo Mt.

Amara (Celia) messae Baliani, det. Hieke (GUÉORGUIEV, 1996, sub *A. (C.) erratica* (Duft.)). A new taxon for Osogovo Mt.

***Amara (Bradytus) consularis* (Duft.)**, det Hieke (GUÉORGUIEV, 1996, sub *A. (B.) fulva* (O. F. Muller)). A first record for Osogovo Mt.

Harpalus (Harpalus) pumilus Sturm, det. Wrase (GUÉORGUIEV, 1996, sub *Egadroma marginata* (Dej.)). *H. pumilus* has been found in both the Bulgarian and the Macedonian parts of Osogovo Mt.

The specimens of the following species, collected from the studied area (GUÉORGUIEV, 1996), belong to the nominate subspecies: *Calosoma auropunctatum* (Hbst.), *Carabus hortensis* L., *Poecilus lepidus* (Leske), *Pterostichus niger* (Schall.), *Pterostichus oblongopunctatus* (F.), *Abax carinatus* (Duft.), *Calathus melanocephalus* (L.), *Calathus erratus* Sahlbg., *Amara apricaria* (Payk.), *Amara equestris* (Duft.), *Gynandromorphus etruscus* (Quens.), *Harpalus rufipalpis* Sturm, *Harpalus quadripunctatus* Dej., and *Harpalus serripes* (Quens.).

Conclusions

As a result of the earlier papers of the author (GUÉORGUIEV, 1996; 1997; 1998), as well as the present one (including the above mentioned corrections) the total

amount of the known ground-beetles taxa from the Osogovo Mt has been increased to 162 species and subspecies. So far 161 from them have been established in the Bulgarian part of the area and 25 in the Macedonian (GUÉORGUIEV, 1998). *Bembidion (Peryphanes) pindicum* Apf. is new for the Bulgarian fauna, while *Pterostichus (Haptoderus) brevis brevis* (Duft.) has been found for the second time in this country.

Acknowledgements

I would like to thank to some colleagues for their help in the determination or confirmation of some specimens: Dr F. Hieke (Berlin) - some *Amara*, Dr B. Kataev (Sankt-Petersburg) - some *Harpalus*, Dr R. Sciaky (Milano) - all *Trechus* and Mr. D. Wräse (Berlin) - various taxa.

References

- APFELBECK V. 1902. Bericht über eine entomologische Forschungsreise nach der Türkei und Griechenland im Jahre 1900. - Wiss. Mitt. Bosn. Hercegov., **8**: 447-469.
- APFELBECK V. 1904. Die Käferfauna der Balkanhalbinsel, mit Berücksichtigung Klein-Asien und der Insel Kreta. Erstes Band: Familinreiche Caraboidea. Berlin, R. Friedlander & Sohn. IX + 422 p.
- BURESCHE I., S. KANTARDJIEVA. 1928. Die in Bulgarien vorkommenden Arten der Subfamilie Carabinae (Coleopt.-Carabidae). - Mitt. K. naturw. Inst. Sofia, **1**: 45-107. (In Bulgarian).
- DRENSKY P. 1928. Referate und Berichte in Jahre 1926-1927. - Mitt. Bulg. ent. Ges. Sofia, **4**: 12-24. (In Bulgarian).
- GUÉORGUIEV B. 1996. A contribution to the study of the ground-beetle fauna (Coleoptera, Carabidae) from the Osogovo Mountain. I. - Hist. nat. bulg., **6**: 29-35.
- GUÉORGUIEV B. 1997. Contribution to the study of the ground-beetle fauna of Osogovo Mountain (Bulgaria). II. Morphological and taxonomic investigations of the genus *Molops* Bonelli (Coleoptera: Carabidae: Pterostichini). - Hist. nat. bulg., **7**: 19-27.
- GUÉORGUIEV B. 1998. Ground-Beetles (Coleoptera: Carabidae) collected of Bulgarian zoologists in Republic of Macedonia. - Hist. nat. bulg., **9**: 35-51.
- GUÉORGUIEV V. 1992. Contribution à l'étude de la famille des Carabidae (Coleoptera) en Bulgarie. II. - Acta zool. bulg., **43**: 61-68. (In Bulgarian).
- HIEKE F., D.W. WRÄSE. 1988. Faunistik der Laufkäfer Bulgariens (Coleoptera, Carabidae). - Dtsch. ent. Z. (N. F.), **35** (1-3): 1-171.
- KANTARDJIEVA S. 1928. Die Arten der Familie Cicindelidae (Col.) in Bulgarien. - Mitt. Bulg. ent. Ges. Sofia, **4**: 91-114. (In Bulgarian).
- MEIXNER J. 1939. Probleme der Rassendifferenzierung, aufgezeigt an Arten der Laufkäfergattung *Trechus*. - In: Jordan K., E. Hering (eds). Verhandlungen. VII. Internationaler Kongress für Entomologie. Band I. Weimar, G. Uschmann, 303-318.

- NETOLITZKY F. 1943. Bestimmungstabelle der *Bembidion*-Arten des paläarktischen Gebiets.
- Koleopt. Rundsch., **29**: 1-70.
- PAWLOWSKI J. 1973. Espèces bulgares du genre *Trechus* (Coleoptera, Carabidae). - Acta Zool. Cracov., **18** (10): 217-270.

Received on 6.7.1998

Author's address:

Dr Borislav Guéorguiev
National Museum of Natural History
1, Tsar Osvoboditel Blvd
1000 Sofia, Bulgaria

Принос към изучаването на бръмбарите-бегачи (Coleoptera: Carabidae) от Осоговската планина. III.

Борислав ГЕОРИГИЕВ

(Резюме)

Настоящата трета поредна работа на автора върху Carabidae от Осоговската планина се състои от две части. В първата са включени 77 нови вида и подвида за планината с данни за вертикалното разпространение, сезонната активност и микроусловията на живот на имагото. *Bembidion (Peryphanes) pindicum* Apf. е нов за българската фауна, а *Pterostichus (Haptoderus) brevis brevis* (Duft.) е намерен за втори път у нас. Материалът е събиран от автора от 29 различни находища през периодите IV-XI.1994, IV-X.1995, IV-X.1996, X.1997 и VI-VII.1998. Включени са и единични екземпляри, събиращи от други лица преди този период, които бяха намерени в колекциите на Националния природонаучен музей в София. Във втората част от работата са коригирани погрешно определените в предишни публикации таксони и е уточнен подвидовият статус на всички публикувани за района видове и подвидове. В резултат общият брой на известните карабиди от Осогово става 162 (161 от българската и 25 от македонската част на планината).